



Product Data Sheet

DMEM High Glucose, With high glucose, L-glutamine, 25 mM HEPES, and sodium pyruvate. Without phenol red and sodium bicarbonate,

Product No. GB-DMEM-10LT, GB-DMEM-50LT

Packing 10 Liters / 50 Liters

Product Description

Appearance	Off-white to creamy white, homogenous powder
pH w/o Sodium Bicarbonate	5.90 – 6.50
pH with Sodium Bicarbonate	7.40 – 8.00
CO ₂ concentration optimum for liquid medium	8.5 %
Storage and shelf life	Store at +2°C to +8°C, dry and protected from light. Please refer to product label for expiration date.
Shipping conditions	Ambient
Use at	13.5 g/L
Add	3.7 g/L Sodium Bicarbonate

Product Description

Preparation of 1 liter liquid medium

1. Suspend 13.5 g in 900 ml tissue culture grade water with constant, gentle stirring until the powder is completely dissolved. Do not heat the water.
2. Add 3.7 g of sodium bicarbonate powder or 49.3 ml of 7.5 % sodium bicarbonate solution for 1 liter of medium and stir until dissolved.
3. Adjust the pH to 0.2 to 0.3 pH units below the desired pH using 1 N HCl or 1 N NaOH since the pH tends to rise during filtration.
4. Add tissue culture grade water up to the final volume of 1000 ml.
5. Sterilize the medium immediately by filtering through a sterile membrane filter with porosity of 0.22 micron or less, using positive pressure rather than vacuum to minimize the loss of carbon dioxide.
6. Aseptically add sterile supplements as required and dispense the desired amount of sterile medium into sterile containers.
7. Store liquid medium at +2°C to +8°C and in dark until use.

Additional Information

- Preparation of concentrated medium is not recommended since free base amino acids and salt complexes having low solubility may precipitate in concentrated medium.



GOSLAR BIOTECH

Your Partner in Sera

- pH and sodium bicarbonate concentration of the prepared medium are critical factors affecting cell growth. This is also influenced by amount of medium and volume of culture vessel used (surface to volume ratio). For example, in large bottles pH tends to rise perceptibly as significant volume of carbon dioxide is released. Therefore, optimal conditions of pH, sodium bicarbonate concentration, surface to volume ratio must be determined for each cell type. We recommend stringent monitoring of pH. If needed, pH can be adjusted by using sterile 1 N HCl or 1 N NaOH or by bubbling in carbon dioxide.

If required, supplements can be added to the medium prior to or after filter sterilization observing sterility precautions. Shelf life of the medium will depend on the nature of supplement added to the medium.

Formulation

Components	Concentration mg/L
Amino Acids:	
L-Arginine HCl	84.00
L-Cystine 2 HCl	62.57
L-Glutamine	584.00
Glycine	30.00
L-Histidine HCl H ₂ O	42.00
L-Isoleucine	105.00
L-Leucine	105.00
L-Lysine HCl	146.00
L-Methionine	30.00
L-Phenylalanine	66.00
L-Serine	42.00
L-Threonine	95.00
L-Tryptophan	16.00
L-Tyrosine 2 Na	103.79
L-Valine	94.00

Components	Concentration mg/L
Vitamins:	
Choline Chloride	4.00
D-Calcium Pantothenate	4.00
Folic Acid	4.00
myo-Inositol	7.20
Nicotinamide	4.00
Pyridoxal HCl	4.00
Riboflavin	0.40
Thiamine HCl	4.00
Inorganic Salts:	
Calcium Chloride 2 H ₂ O	265.00
Fe(NO ₃) ₃ 9 H ₂ O	0.10
MgSO ₄ (Anhydrous)	97.72
KCl	400.00
NaCl	6400.00
NaH ₂ PO ₄ (Anhydrous)	109.00
Other Components:	
D-Glucose	4500.00
Phenol Red Sodium Salt	15.90
Sodium Pyruvate	110.00

Precautions and Disclaimer

This product is for research use only.