

Product Data Sheet

ACTINOMYCETE ISOLATION AGAR

Product No. GB-DCM-00014-1A

INTENDED USE

For isolation and propagation of Actinomycetes from soil and water.

Product Description

Actinomycetes are gram-positive bacteria, which show marked chemical and morphological diversity but form a distinct evolutionary line of organisms that range from coccoid and pleomorphic forms to branched filaments. Actinomycetes form an integral part of soil, water and vegetation. Actinomycete development leads to the formation of volatile metabolites. Traces of these volatile metabolites are sufficient to impart disagreeable odour to water or a muddy flavour to fish. Actinomycetes also cause disruptions in wastewater treatment by forming massive growths, which are capable of producing thick foam in the activated sludge process. Actinomycetes Isolation Agar used for isolation and propagation of Actinomycetes from soil and water was formulated by Olsen. Inoculate the plates with 1 drop of diluted culture or specimen and spread over the surface using a sterile bent glass rod. Incubate at 35-37°C for 40-72 hours. The media can be used for long term storage after sufficient growth is obtained. Agar slants are used for maintenance of cultures over a shorter period of time.

Product Specifications

Ingredients	Gms / Ltr
Sodium caseinate	2.000
L-Asparagine	0.100
Sodium propionate	4.000
Dipotassium phosphate	0.500
Magnesium sulphate	0.100
Ferrous sulphate	0.001
Agar	15.000

PRINCIPLE

Actinomycete Isolation Agar contains sodium caseinate as nitrogen source. Asparagine in addition to being an amino acid is also a source of nitrogen. Sodium propionate is used as a substrate in anaerobic fermentation. Dipotassium phosphate provides the buffering system. The sulphates serve as source of sulphur and metallic ions. Glycerol serves as an additional source of carbon.



INSTRUCTION FOR USE

- Dissolve 21.70 grams in 1000 ml distilled water containing 5 ml glycerol.
- Heat to boiling to dissolve the medium completely. Dispense as desired.
- Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder: Cream to yellow homogeneous free flowing powder. Appearance of prepared medium: Yellow to light amber coloured opalescent gel forms in Petri plates.
pH (at 25°C) : 8.1±0.2

Microorganism	ATCC	Inoculum (CFU)	Growth	Recovery	Incubation Temperature	Incubation Period
Nocardia asteroides	19427	50-100	Good-luxuriant	≥50%	35-37°C	40-72 Hours
Escherichia coli	25922	≥104	Inhibited	0%	35-37°C	40-72 Hours
Streptomyces albus subsp albus	3004	50-100	Good-luxuriant	≥50%	35-37°C	40-72 Hours
Streptomyces lavendulae	19247	50-100	Good-luxuriant	≥50%	35-37°C	40-72 Hours

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

This product is for research use only.