



Product Data Sheet

ASHBY'S GLUCOSE AGAR

Product No. GB-DCM-00045-1A

INTENDED USE

For cultivation of Azotobacter species by using glucose as carbon source.

Product Description

Azotobacter is a genus of free-living diazotrophic bacteria which have the highest metabolic rate compared to any other microorganism. Azotobacters are chemoorganotrophic, using sugars, alcohols and salts of organic acids for growth. Ashby's Agar Media are formulated as described by Subba Rao. It is used for isolation of Azotobacter, a non-symbiotic nitrogen fixing bacteria which uses glucose as a carbon source and atmospheric nitrogen as nitrogen source. Besides the ability to fix atmospheric nitrogen, Azotobacter also synthesizes biologically active substances which attributes to improving seed germination, plant growth etc.

Product Specifications

Ingredients	Gms / Ltr
Glucose (Glucose)	20.000
Dipotassium hydrogen phosphate	0.200
Magnesium sulphate	0.200
Sodium chloride	0.200
Potassium sulphate	0.100
Calcium carbonate	5.000
Agar	15.000

PRINCIPLE

Dipotassium phosphate provides buffering to the system. Various essential ions required for promoting growth of Azotobacter are also available in this medium.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder: Cream to yellow homogeneous free flowing powder.

Appearance of prepared medium: Whitish, opalescent gel forms in Petri plates.

PH (at 25°C): 7.4±0.2

Microorganism	ATCC	Inoculum (CFU)	Growth	Recovery	Incubation Temperature	Incubation Period
Azotobacter vinelandii	478	50-100	Good-Luxuriant	>=50%	35-37°C	5 Days

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration:

Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

Precautions and Disclaimer

- Suspend 40.7 grams in 1000 ml purified/distilled water.
- Heat if necessary to dissolve the medium completely.
- Mix well and pour into sterile Petri plates.
- Sterilize by autoclaving at 15 psi pressure (121°C) for 10 minutes.

This product is for research use only.