

## Product Data Sheet

### CHOCOLATE AGAR BASE

**Product No.** GB-DCM-00070-1A

### INTENDED USE

For differentiation of enteric pathogens and coliforms on the basis of citrate utilization.

### PRODUCT SUMMARY

Christensen Citrate Agar is a modification of Christensen Iron Agar, which has the same formulation except the additional sodium thiosulphate and ferric ammonium citrate as described by Edwards and Ewing. Christensen reported that all members of genera Escherichia, Enterobacter, Citrobacter and Salmonella were capable of utilizing citrate as a source of energy while Shigella species failed to utilize citrate. Edward and Ewing recommended the use of Triple Sugar Iron Agar for the determination of hydrogen sulphide production and Christensen Citrate Agar for citrate utilization. Organisms that metabolize citrate as a sole source of carbon cleave citrate to oxaloacetate and acetate via the citritase enzyme. Another enzyme, oxaloacetate decarboxylase, then converts oxaloacetate to pyruvate and CO<sub>2</sub>. Further, this CO<sub>2</sub> combines with sodium and water to form sodium carbonate, an alkaline compound. As a result, the pH of medium rises and the indicator, phenol red changes from orange red to cerise. Presence of the cerise colour indicates a positive finding for citrate utilization.

### Product Specifications

Ingredients	Gms / Ltr
Yeast extract 0.500	0.500
L-Cysteine hydrochloride 0.100	0.100
Sodium citrate 3.000	3.000
Dextrose	0.200
Potassium dihydrogen phosphate 1.000	1.000
Sodium chloride	5.000
Sodium chloride	0.012
Agar	15.000

### PRINCIPLE

Medium constituent yeast extract provide the necessary nutrients mainly nitrogenous and vitamins for the growth of the organisms. L-cysteine hydrochloride is a reducing agent. Dextrose is the fermentable carbohydrate. Sodium citrate is the energy source for citrate utilizing organisms.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Recovery	Citrate utilization (colour of slant)	Incubation Temperature	Incubation Period
Klebsiella aerogenes	13048	50-100	luxuriant	>=70%	Positive reaction, cerise colour	35-37°C	24-48 Hours
Escherichia coli	13883	50-100	luxuriant	>=70%	Negative reaction, no colour change	35-37°C	24-48 Hours
Klebsiella pneumoniae	13883	50-100	luxuriant	>=70%	Weakly positive, orange-pink colour	35-37°C	24-48 Hours
Salmonella Typhimurium	14028	50-100	luxuriant	>=70%	Positive reaction, cerise colour	35-37°C	24-48 Hours
Salmonella Enteritidis	13076	50-100	luxuriant	>=70%	Positive reaction, cerise colour	35-37°C	24-48 Hours
Shigella flexneri	12022	50-100	luxuriant	>=70%	Negative reaction, no colour change	35-37°C	24-48 Hours
Shigella sonnei	25931	50-100	luxuriant	>=70%	Negative reaction, no colour change	35-37°C	24-48 Hours

### INSTRUCTION FOR USE

Dissolve 24.8 grams in 1000 ml purified / distilled water.

- Heat to boiling to dissolve the medium completely.
- Dispense into test tubes. Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.
- Cool the tubes in a slanted position.

### QUALITY CONTROL SPECIFICATIONS

Appearance of Powder.

Light yellow to light pink homogeneous free flowing powder.

Appearance of prepared medium: tubes as slants.

Orange red coloured, very slightly opalescent gel forms in

pH (at 25°C) : 6.9±0.2



**GOSLAR BIOTECH**  
*Your Partner in Sera*

### **STORAGE**

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

**Product Deterioration:** Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

### **DISPOSAL**

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

**This product is for research use only.**