

Product Data Sheet

CAL AGAR (CELLOBIOSE ARGININE LYSINE AGAR)

Product No. GB-DCM-00140-1A

INTENDED USE

For isolation and biochemical identification of *Yersinia enterocolitica*.

Product Description

Yersinia enterocolitica is a significant invasive enteric pathogen belonging to the family Enterobacteriaceae, which causes several well-recognized diseases especially in younger persons and several uncommon post-infection syndromes. Enterocolitis caused by *Y. enterocolitica* is characterized by diarrhoea, low fever and abdominal pain. CAL Agar used for selective isolation of *Y. enterocolitica* was originally formulated by Dudley and Shotts. CAL Agar is a differential medium as it differentiates *Yersinia* on the basis of cellobiose fermentation and lysine or arginine decarboxylation. CAL Agar is generally used for the isolation and characterization of *Y. enterocolitica* from faecal specimens as the organism is biochemically similar to other Enterobacteriaceae. CAL Broth is used for the enumeration of *Y. enterocolitica* from water and other liquid specimens.

COMPOSITION

Ingredients	Gms / Ltr
Yeast extract	3.000
Sodium chloride	5.000
Cellobiose	3.500
L-Arginine	6.500
L-Lysine hydrochloride	6.500
Sodium deoxycholate	1.500
Neutral red	0.030
Agar	20.000

PRINCIPLE

Yeast extract provides essential nutrients to the organisms. Cellobiose is the fermentable carbohydrate. Sodium chloride maintains the osmotic equilibrium. Sodium deoxycholate makes the medium selective by inhibiting the accompanying gram-positive bacteria, which may cause contamination during cultivation. L-arginine and L-lysine are the amino acids, decarboxylation of which makes the medium differential. Neutral red is the indicator, which turns red under acidic conditions.

INSTRUCTION FOR USE

- Dissolve 46.03 grams in 1000 ml distilled water.
- Heat to boiling to dissolve the medium completely.
- Cool to 45-50°C. Mix well and pour into sterile Petri plates.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder: Cream to yellow homogeneous free flowing powder.
 Appearance of prepared medium: Red coloured, clear to slightly opalescent gel forms in Petri plates.
 pH (at 25°C): 7.1±0.2

Microorganism	ATCC	Inoculum (CFU)	Growth	Recovery	Cellobiose	Arginine Decarboxylation	Lysine Decarboxylation	Incubation Temperature	Incubation Period
Candida albicans	25922	50-100	Good	40- 50%	Negative reaction	Variable reaction	Variable reaction	35-37°C	18-48 Hours
Aspergillus brasiliensis	25933	50-100	Good	40- 50%	Negative reaction	Negative reaction	Negative reaction	35-37°C	18-48 Hours
Candida albicans	27853	50-100	Good	40- 50%	Negative reaction	Negative reaction	Positive reaction	35-37°C	18-48 Hours
Saccharomyces cerevisiae	27729	50-100	Good-luxuriant	>=50%	Positive reaction	Negative reaction	Negative reaction	35-37°C	18-48 Hours



STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

This product is for research use only.