



Product Data Sheet

COLUMBIA AGAR (as per USP/EP/JP/BP/IP)

Product No. GB-DCM-00147-1A

INTENDED USE

For detection of Clostridium sporogenes from pharmaceutical products.

PRODUCT SUMMARY

Columbia Blood Agar Base used as a general-purpose nutritious medium was devised by Elmer al from Columbia University, which was further enriched by the addition of sheep blood. It can also be used for the isolation of organisms by addition of various supplements. Columbia Agar is prepared in accordance with the microbial limit testing harmonized methodology of USP/EP/BP/JP/IP. This medium is recommended to check the presence of Clostridium spp. in non-sterile products like food, dietary, nutritional supplements related products. Columbia agar base can be used to prepare lactose milk egg-yolk agar for the isolation of fastidious Clostridia. Al-Jumaili and Bint (1981) recommended the addition of blood, cycloserine and cefoxitin to Columbia agar (base) for the isolation of Clostridium difficile. The inclusion of bacitracin makes the enriched Columbia Agar Medium selective for the isolation of Haemophilusspecies from clinical specimens, especially from upper respiratory tract.

Product Specifications

Ingredients	Gms / Ltr
Agar	15.000
Pancreatic digest of Casein	10.000
Peptic digest of meat	5.000
Yeast extract	5.000
Sodium chloride	5.000
Pancreatic digest of heart	3.000
Maize starch	1.000

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

INSTRUCTION FOR USE

- Dissolve 44.00 grams in 1000 ml distilled water.
- Gently heat to boiling with swirling to dissolve the medium completely.
- Sterilize by autoclaving at psi (121°C) for 15 minutes.
- Cool to 45-50 °C and if required add the rehydrated contents of 1 vial of Gentamycin supplement (TS 217).
- Mix well and pour into sterile petri

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Recovery	Incubation Temperature	Incubation Period
Clostridium sporogenes	11437	50-100	Good - Luxuriant	≥50%	35-37°C	≤48 hours
Clostridium sporogenes	19404	50-100	Good - Luxuriant	≥50%	35-37°C	≤48 hours
Clostridium perfringens	13124	50-100	Good - Luxuriant	≥50%	35-37°C	≤48 hours
Bacteroides fragilis	23745	50-100	Good - Luxuriant	≥50%	35-37°C	≤48 hours

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder. Cream to yellow homogeneous free flowing powder.
 Appearance of prepared medium: Light amber colour, opalescent gel.
 pH (at 25°C) : 7.3±0.2

PRINCIPLE

This medium is highly nutritious as it contains Pancreatic digest of Casein, Peptic digest of meat, Pancreatic digest of heart and Yeast extract, which provides carbonaceous and nitrogenous substances, long chain amino acids, vitamins of B complex group and other essential nutrients for the luxuriant growth of fastidious as well as non-fastidious organisms. Sodium chloride maintains osmotic balance of medium. Maize starch acts as an energy source and also neutralizes toxic metabolites if produced. It is used in detection of Clostridia from pharmaceutical products. Agar acts as a solidifying agent. Gentamycin supplement (TS 217), when added acts as a selective agent against a number of gram negative organisms and also Staphylococcus species.



GOSLAR BIOTECH
Your Partner in Sera

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

This product is for research use only.