

Product Data Sheet

HEART INFUSION BROTH (BEEF HEART INFUSION BROTH)

Product No. GB-DCM-00250-1A

INTENDED USE

For isolation and cultivation of various fastidious microorganisms.

PRODUCT SUMMARY

Fastidious organisms having exacting nutritional requirement could be cultivated on infusion media, as demonstrated by Huntoon. A liquid medium containing an infusion of meat was one of the first media used for the cultivation of bacteria. These infusion media need not be further supplemented by the addition of supplements for cultivation of fastidious bacteria. HI Agar, containing infusion from beef heart is used for the isolation and cultivation of a wide variety of fastidious organisms. HI Agar can also be used for the cultivation of *Vibrio* species. It can also be supplemented with glucose, horse serum and antibiotics for the cultivation a wide variety of organisms. It is used for mass cultivation of organisms for preparation of vaccines. On supplementation of blood, HI Agar can be used to study haemolytic reactions. This medium was used for isolation and enumeration of haemolytic *Streptococci* in milk.

Product Specifications

Ingredients	Gms / Ltr
Beef heart, infusion from	500.00
Tryptose	10.000
Sodium chloride	5.000

PRINCIPLE

Tryptose and Beef heart, infusion from provide nutritional requirements for the pathogenic bacteria. Sodium chloride maintains the osmotic equilibrium of the medium.

INSTRUCTION FOR USE

- Dissolve 25.0 grams in 1000 ml of distilled water.
- Heat if necessary to dissolve the medium completely.
- Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.
- If desired 5% v/v sterile defibrinated blood may be added.
- Mix well and aseptically pour into sterile tubes or flasks as desired.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.



QUALITY CONTROL SPECIFICATIONS

Appearance of Powder: Cream to yellow homogeneous free flowing powder
 Appearance of prepared medium: Basal medium: Light yellow coloured, clear to slightly opalescent solution. After addition of 5% sterile defibrinated blood: Cherry red coloured, opaque solution in tubes.
 pH (at 25°C) : 7.4 ± 0.2

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Incubation Temperature	Incubation Period
Escherichia coli	25922	50-100	Good-Luxuriant	35-37°C	18-48 Hours
Neisseria meningitidis	13090	50-100	Good-Luxuriant	35-37°C	18-48 Hours
Streptococcus pneumoniae	6303	50-100	Good-Luxuriant	35-37°C	18-48 Hours
Streptococcus pyogenes	19615	50-100	Good-Luxuriant	35-37°C	18-48 Hours
Staphylococcus aureus subsp. aureus	25923	50-100	Good-Luxuriant	35-37°C	18-48 Hours

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

This product is for research use only.