



## **Product Data Sheet**

### **HIGH SENSITIVITY TEST BROTH**

**Product No.** GB-DCM-00253-1A

### **INTENDED USE**

For antimicrobial susceptibility tests.

### **PRODUCT SUMMARY**

The goal of an antimicrobial susceptibility test is to predict through an in vitro assessment the likelihood of successfully treating an infection with a particular antimicrobial agent. There are several continual or novel methods for performing antibacterial susceptibility testing. These include the disk diffusion test, broth microdilution, agar gradient and rapid automated instrument methods. Hi-Sensitivity Test Broth, which is used for antimicrobial susceptibility tests, is a semidefined medium in which the mineral contents have been stabilized to give reproducible results. The thiamine and thymidine content is very low thus making it most suitable for testing antimicrobial activity of sulphonamides. However, some mutant strains which are totally dependent on thiamine and thymidine for their growth, will not grow in HiSensitivity Test Broth, due to very low levels of these compounds in the media as they are the naturally occurring antagonist of trimethoprim. These strains should be carefully recognized. Hi-Sensitivity Test Broth has been so designed to overcome the problems occurring in Mueller-Hinton Media that are as follows. 1. Mueller Hinton Agar and Mueller Hinton Broth give different MIC values. 2. Mueller Hinton Agar shows antagonistic effect towards tetracycline. 3. High levels of sulphonamide and trimethoprim antagonists. 4. Media prepared using peptone of different manufacturers give poor reproducibility. 5. Poor growth supporting ability for Streptococci and variable growth rates with gram-positive organisms. Some pathogenic organisms are nutritionally dependent due to their intrinsic demands for special growth factors.

### **PRINCIPLE**

Casein enzymic hydrolysate, peptic digest of animal tissue, dextrose, and vitamins provides nitrogen, carbon compounds and other essential growth nutrients.

### **INSTRUCTION FOR USE**

- Dissolve 23.40 grams in 1000 ml distilled water.
- Heat to boiling to dissolve the medium completely.
- Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.
- Mix well and dispense as desired.



**GOSLAR BIOTECH**  
*Your Partner in Sera*

## Product Specifications

<b>Ingredients</b>	<b>Gms / Ltr</b>
Casein enzymic hydrolysate	11.000
Peptic digest of animal tissue	3.000
Dextrose	2.000
Sodium chloride	3.000
Starch, soluble	1.000
Disodium phosphate	2.000
Sodium acetate	1.000
Magnesium glycerophosphate	0.200
Calcium gluconate	0.100
Cobaltous sulphate	0.001
Cupric sulphate	0.001
Ferrous sulphate	0.001
Zinc sulphate	0.001
Manganous chloride	0.002
Menadione	0.002
Cyanocobalamin	0.002
L-Cysteine hydrochloride	0.020
L-Tryptophan	0.020
Pyridoxine hydrochloride	0.003
Calcium pantothenate	0.003
Nicotinamide	0.003
Biotin 0.0003	0.0003
Thiamine hydrochloride	0.00004
Adenine	0.010
Guanine	0.010
Xanthine	0.010
Uracil	0.010



## QUALITY CONTROL SPECIFICATIONS

Appearance of Powder: Cream to yellow homogeneous free flowing powder  
 Appearance of prepared medium: Basal medium: Light yellow; After addition of 5%v/v laked blood: Red to chocolate coloured, Basal medium: clear to slightly opalescent; After Addition: opalescent solution in tubes.  
 pH (at 25°C) : 7.4 ± 0.2

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Incubation Temperature	Incubation Period
Salmonella Typhimurium	14028	50-100	Good-Luxuriant	35-37°C	18-24 Hours
Staphylococcus aureus	25923	50-100	Good-Luxuriant	35-37°C	18-24 Hours
Streptococcus pyogenes	19615	50-100	Good-Luxuriant	35-37°C	18-24 Hours
Enterococcus faecalis	29212	50-100	Good-Luxuriant	35-37°C	18-24 Hours
Bacillus subtilis	6633	50-100	Good-Luxuriant	35-37°C	18-24 Hours
Bacteroides vulgatus	8482	50-100	Good-Luxuriant	35-37°C	18-24 Hours

## STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

**Product Deterioration:** Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

## DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

**This product is for research use only.**