

Product Data Sheet

LACTOBACILLI BROTH (AOAC)

Product No. GB-DCM-00296-1A

INTENDED USE

For preparation of inocula of stock cultures used in microbiological assays of vitamins of group B.

PRODUCT SUMMARY

Lactobacilli Broth, AOAC was formulated by Loy and recommended by AOAC for preparing inocula of test bacteria used for Microbiological assay of Vitamin B. Stock cultures of Lactobacillus leichmanni ATCC 7830, Lactobacillus plantarum ATCC 8014, Lactobacillus casei ATCC 7469, Enterococcus hirae ATCC 8043 and other such B vitamin requiring strains are prepared by stab inoculation of sterile Lactobacillus Agar, AOAC and incubated for 18-24 hours at a constant temperature between 30-40°C. Lactobacilli Broth, AOAC is used for cultivation and preparation of inocula of the above mentioned stock cultures. Inoculum is prepared by inoculating these cultures in Lactobacillus Broth, AOAC and incubating at 35-37°C.

Product Specifications

Ingredients	Gms / Ltr
Peptonized milk	15.000
Yeast extract	5.000
Dextrose (Glucose)	10.000
Tomato juice (100 ml)	5.000
Potassium dihydrogen phosphate	2.000
Polysorbate 80 (Tween 80)	1.000

PRINCIPLE

This medium consists of Peptonized milk and yeast extract which provide essential growth nutrients. Dextrose is the energy source. Phosphate provides buffering system while tomato juice helps in lowering the pH. Polysorbate 80 supplies fatty acids. Before using a culture in any assay, at least 2 successive transfers during a 1-2 week period are essential. Any culture older than one week should not be used.



GOSLAR BIOTECH
Your Partner in Sera

INSTRUCTION FOR USE

- Dissolve 38.0 grams in 1000 ml purified/distilled water.
- Heat if necessary to dissolve the medium completely.
- Distribute into tubes in 10 ml amounts and sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder : Cream to yellow homogeneous free flowing powder.

Appearance of prepared medium Medium amber coloured clear solution in tubes.

pH (at 25°C) : 6.8± 0.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Incubation Temperature	Incubation Period
Enterococcus hirae	8043	50-100	Luxuriant	35-37 °C	18-48 Hours
Lactobacillus casei	7469	50-100	Luxuriant	35-37 °C	18-48 Hours
Lactobacillus leichmannii	7830	50-100	Luxuriant	35-37 °C	18-48 Hours
Lactobacillus plantarum	8014	50-100	Luxuriant	35-37 °C	18-48 Hours

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

This product is for research use only.

Goslar Biotech, 255A Barking Road East Ham, London E6 1LB, United Kingdom
Email: info@goslarbiotech.com, Website: www.goslarbiotech.com