

Product Data Sheet

LACTIC BACTERIA DIFFERENTIAL BROTH

Product No. GB-DCM-00300-1A

INTENDED USE

For differentiation of homo-fermentative and hetero-fermentative lactic acid bacteria.

PRODUCT SUMMARY

Lactic Bacteria Differential Broth is formulated as per McDonald et al for differentiation of homofermentative Lactobacilli and heterofermentative Streptococci. Lactobacilli and Streptococci are used as starter cultures in food and dairy industry. Streptococci grow first and produce metabolites, lowering redox potential which enables Lactobacilli to grow. Lactobacilli synthesize products which stimulate growth of Streptococci. Heterofermentative lactic acid bacteria produce CO₂, lactic acid, acetic acid, ethanol and mannitol. Homofermentative lactic acid bacteria produce only lactic acid. Homofermentative lactic acid bacteria produce lactic acid from fructose and is indicated by yellow colour formation. Heterofermentative lactic acid bacteria induce lesser acidification and thus vary in the colour formation by the indicator in the medium.

Product Specifications

Ingredients	Gms / Ltr
Tryptone	10.000
Soya peptone	1.500
Casein acid hydrolysate	3.000
Yeast extract	1.000
Fructose	2.500
Potassium dihydrogen phosphate	2.500
Bromocresol green	0.055

PRINCIPLE

This medium consists of Casein acid hydrolysate, soya peptone and yeast extract which supplies nitrogen and carbon compounds, long chain amino acids, vitamins and all the necessary nutrients for the growth of lactic bacteria. Fructose is the fermentable carbohydrate in the medium. Bromo cresol green is the pH indicator.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.



GOSLAR BIOTECH
Your Partner in Sera

INSTRUCTION FOR USE

- Dissolve 20.55 grams in 1000 ml purified/distilled water. Add 1 gram of polysorbate 80.
- Heat if necessary to dissolve the medium completely.
- Dispense as desired and Sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder : Light yellow to bluish grey homogeneous free flowing powder.

Appearance of prepared medium: Blue coloured clear solution in tubes.

pH (at 25°C) : 7.0± 0.2

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Colour of medium	Incubation Temperature	Incubation Period
Lactobacillus casei	9595	50-100	Luxuriant	Green	35-37°C	18-48 Hours
Lactobacillus plantarum	8014	50-100	Luxuriant	Green	35-37°C	18-48 Hours
Streptococcus thermophilus	14485	50-100	Luxuriant (incubated at 45°C)	Bluish-green	35-37°C	18-48 Hours
Streptococcus cremoris	19257	50-100	Luxuriant (incubated at 30°C)	Blue	35-37°C	18-48 Hours

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

This product is for research use only.

Goslar Biotech, 255A Barking Road East Ham, London E6 1LB, United Kingdom
Email: info@goslarbiotech.com, Website: www.goslarbiotech.com