

Product Data Sheet

MIU MEDIUM BASE
Product No. GB-DCM-00362-1A

INTENDED USE

For detection of motility, urease and indole production.

PRODUCT SUMMARY

MIU Medium Base is formulated to detect motility, urease and indole production in single tube. Motility and urease reactions are read before testing Indole production. Motile organisms show either diffused growth or turbidity extending away from stab inoculation line while nonmotile organisms grow along the stabline. Organisms that utilize urea, produce ammonia which makes the medium alkaline, showing pink-red colour by change in the phenol red indicator. Indole is produced from tryptophan present in tryptone. The indole produced combines with the aldehyde present in the Kovac's reagent to form a red complex.

Product Specifications

Ingredients	Gms / Ltr			
Tryptone	10.000			
Dextrose (Glucose)	1.000			
Sodium chloride	5.000			
Phenol red	0.010			
Agar	2.000			

PRINCIPLE

Tryptone provide amino acids and other nitrogenous substances. Sodium chloride maintains osmotic equilibrium. Dextrose is fermentable carbohydrate. Phenol red is the pH indicator which turns pink- red in alkaline conditions. The test cultures are stab-inoculated.

INSTRUCTION FOR USE

- Dissolve 18 grams in 950 ml purified / distilled water. Heat to boiling to dissolve the medium completely.
- Dispense in 95 ml amounts into flasks and sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.
- Cool to about 50-55°C and add aseptically 5 ml sterile 40% Urea solution per 95 ml basal medium.
- Mix well and dispense into sterile test tubes.
- Allow to cool in an upright position.



UALITY CONTROL SPECIFICATIONS

Appearance of Powder: Light orange to light pink coloured homogeneous free flowing powder. Appearance of prepared medium: Yellowish orange coloured clear to slightly opalescent gel is obtained in tubes as butts afteraddition of urea solution.

pH (at 25°C): 6.8 ± 0.2

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Recovery	Indole	Motility	Urease activity	Incubation Temperature	Incubation Period
Escherichia coli	25922	50-100	luxuriant	>=70 %	Positive reaction, red ring at the interface of the medium	Positive reaction, red ring at the interface of the medium	Negative reaction,no change	35-37°C	18-24 Hours
Klebsiella pneumoniae	13883	50-100	luxuriant	>=70 %	Negative reaction no colour development / cloudy ring	Negative reaction no colour development / cloudy ring	Weakly positive	35-37°C	18-24 Hours
Proteus mirabilis	25933	50-100	luxuriant	>=70 %	Negative reaction no colour development / cloudy ring	Positive reaction, red ring at the interface of the medium	Positive reaction, cerise colour	35-37°C	18-24 Hours
Proteus vulgaris	13315	50-100	luxuriant	>=70 %	Positive reaction, red ring at the interface of the medium	Positive reaction, red ring at the interface of the medium	Positive reaction, cerise colour	35-37°C	18-24 Hours
Salmonella Typhimurium	14028	50-100	luxuriant	>=70 %	Negative reaction no colour development / cloudy ring	Positive reaction, red ring at the interface of the medium	Negative reaction,no change	35-37°C	18-24 Hours

STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

This product is for research use only.

Goslar Biotech, 255A Barking Road East Ham, London E6 1LB, United Kingdom Email: info@goslarbiotech.com, Website: www.goslarbiotech.com