

Product Data Sheet

MANNITOL SALT BROTH (VEG.)

Product No. GB-DCM-00415-1A

INTENDED USE

For selective isolation of presumptive pathogenic Staphylococci.

PRODUCT SUMMARY

These media are prepared by completely replacing animal based peptones with vegetable peptones which makes the medium free of BSE/TSE risks. Mannitol Salt Veg media are the modification of Mannitol Salt media which are prepared as suggested by Chapman and are used for the selective isolation of pathogenic Staphylococci and also are recommended for the detection and enumeration of coagulase-positive Staphylococci in milk food and other specimens.

Product Specifications

Ingredients	Gms / Ltr
Veg peptone No. 3	10.00
Veg extract	1.00
Sodium chloride	75.00
D-Mannitol	10.00
Phenol red	0.025

PRINCIPLE

The medium contains Veg extract and Veg Peptone No. 3 which makes it very nutritious as they provide essential growth factors and trace nutrients. Many other bacteria except Staphylococci are inhibited by 7.5% sodium chloride. Mannitol is the fermentable carbohydrate source. The differential action of the medium is attributed to DMannitol. Staphylococcus aureus grows on this medium and ferment mannitol to produce yellow colonies with yellow zones. Most coagulase-negative species of Staphylococci and Micrococci do not ferment mannitol and grow as small red colonies surrounded by red or purple zones. The colour of colonies and medium is due to the reactivity of phenol red to the pH of the medium; phenol red is red at pH 8.4 and yellow at 6.8. Yellow colonies should be tested for production of coagulase. Addition of 5% v/v Egg Yolk Emulsion enables to detect lipase activity of Staphylococci along with mannitol fermentation. The salt clears egg yolk emulsion and the lipase production is detected as yellow opaque zone around the colonies. Presumptive coagulase-positive Staphylococci produces colonies surrounded by bright yellow zones while non- pathogenic Staphylococci produce colonies with reddish purple zones.

INSTRUCTION FOR USE

- Dissolve 96.02 grams in 1000 ml purified / distilled water.
- Heat if necessary to dissolve the medium completely.
- Dispense as desired and sterilize by autoclaving at 15 psi pressure (121°C) for 15 minutes.
- Cool to 45-50°C. Mix well and dispense as desired.

Microorganism	ATCC	Inoculum (CFU/ml)	Growth	Colour of medium	Incubation Temperature	Incubation Period
Escherichia coli	25922	$\geq 10^4$	Inhibited	Pink with bile precipitate	35-37°C	18-24 Hours
Staphylococcus subsp. aureus	25923	50-100	Good-luxuriant	Colourless	35-37°C	18-24 Hours
Staphylococcus Epidermidis	12228	50-100	Fair-good	Colourless	35-37°C	18-24 Hours
Enterobacter aerogenes	13048	50-100	Inhibited	Colourless	35-37°C	18-24 Hours
Proteus mirabilis	12453	50-100	None to poor	Red	35-37°C	18-24 Hours

QUALITY CONTROL SPECIFICATIONS

Appearance of Powder :	Light yellow to pink homogeneous free flowing powder.
Appearance of prepared medium :	Red coloured, clear to slightly opalescent gel forms in petri Plates, clear solution in tubes.
pH (at 25°C) :	7.4± 0.2



STORAGE

Dehydrated powder, hygroscopic in nature, store in a dry place, in tightly-sealed containers between 25-30°C and protect from direct sunlight. Under optimal conditions, the medium has a shelf life of 4 years. When the container is opened for the first time, note the time and date on the label space provided on the container. After the desired amount of medium has been taken out replace the cap tightly to protect from hydration.

Product Deterioration: Do not use if they show evidence of microbial contamination, discoloration, drying or any other signs of deterioration.

DISPOSAL

After use, prepared plates, specimen/sample containers and other contaminated materials must be sterilized before discarding.

This product is for research use only.